

B.Tech I Year (R13) Supplementary Examinations December/January 2014/2015

MATHEMATICS – II

(Common to EEE, ECE, EIE, CSE and IT)

Time: 3 hours

Max. Marks: 70

PART – A

(Compulsory Question)

1 Answer the following: (10 X 02 = 20 Marks)

- Find the sine series of $f(x) = k$ in $(0, \pi)$.
- If $f(x) = x + x^2$ in $-\pi < x < \pi$ then find a_n .
- Obtain the complete solution for $p + q = \sin x + \sin y$.
- Find $a_0, f(x) = |\cos x|, (-\pi, \pi)$.
- Find P.I of $(D^2 - 2DD') z = x^3 y$.
- State one dimensional heat equation.
- Find the Eigen values for the matrix $\begin{bmatrix} 4 & 2 & -2 \\ -5 & 3 & 2 \\ -2 & 4 & 1 \end{bmatrix}$.
- Write condition for the system $AX = B$ is consistent.
- Find the rank of $\begin{bmatrix} 1 & -9 & 6 \\ 4 & 8 & 5 \\ 7 & 9 & 4 \end{bmatrix}$.
- Using Euler's method find the solution of the initial problem $\frac{dy}{dx} = \log(x + y), y(0) = 2$ at $x = 0.2$ by assuming $h = 0.2$.

PART – B

(Answer all five units, 5 X 10 = 50 Marks)

UNIT - I

- 2 Reduce the quadratic form $3x^2 + 5y^2 + 3z^2 - 2yz + 2zx - 2xy$ to the canonical form. Also specify the matrix of transformation.

OR

- 3 State and prove Cayley-Hamilton theorem.

UNIT - II

- 4 Find the root of $x \log_{10} x - 1.2 = 0$ by Newton Raphson method corrected to three decimal places.

OR

- 5 Evaluate $\int_0^1 x e^x dx$ taking 4 intervals. Using (i) Trapezoidal rule. (ii) Simpson's 1/3 rd rule.

UNIT - III

- 6 Use fourth order Runge-Kutta method to compare y for $x = 0.1$, given $\frac{dy}{dx} = \frac{xy}{1+x^2}, y(0) = 1$ take $h = 0.1$.

OR

- 7 Find the Half range Fourier sine series $f(x) = x(\pi - x) \quad 0 \leq x \leq \pi$ and hence deduce that: (i) $\sum \frac{1}{n^4} = \frac{\pi^4}{960}$.

$$(ii) \sum_{n=1}^{\infty} \frac{1}{(2n-1)^6} = \frac{\pi^6}{960}.$$

UNIT - IV

- 8 Find the Fourier cosine transform of $f(x) = e^{-x^2}$.

OR

- 9 Solve Z-transform $y_{k+1} + \frac{1}{4}y_k = \left(\frac{1}{4}\right)^k, (k \geq 0), y(0) = 0$.

UNIT - V

- 10 Solve the equation $\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}$ with boundary conditions $u(x, 0) = 3 \sin(n\pi x), u(x, t) = 0, u(a, t) = 0$, where $0 < x < 1, t > 0$.

www.jntuonline.com

- 11 A tightly stretched string with fixed end points $x = 0$ and $x = 1$ is initially in a position given by $y = y_0 \sin^3(\pi x/l)$. If it is released from rest from this position, find the displacement $y(x, t)$.
